



CONNECTICUT
KIDNEY & HYPERTENSION
SPECIALISTS

www.ctkidneyspecialists.com

VACCINATION GUIDE / LOG

for PEOPLE with KIDNEY DISEASE

RECOMMENDED VACCINATION FOR ADULTS WITH CHRONIC KIDNEY DISEASE STAGE 4-5, ON DIALYSIS OR WITH A KIDNEY TRANSPLANT					
VACCINE	PT WITH CKD stage 4-5	PT ON DIALYSIS	PT WITH A TRANSPLANT	DATE ADMINISTERED	NOTES
INFLUENZA (Flu)	Recommended	Recommended	Recommended Consider repeat dose 4-8 weeks		Yearly- Quadrivalent (High dose) if age > 65 yo
COVID 19 (Corona virus)	Recommended	Recommended	Recommended		Per CDC recommendations
PNEUMOCOCCAL P13 and P23 or PCV20* (also known as Pneumonia Vaccine)	Recommended	Recommended	Recommended		Repeat dosing at age 65, if given prior
Tetanus-Diphtheria-Pertussis (whooping cough)	Recommended if not done last 10 years	Recommended if not done in last 10 years	Recommended if not done in last 10 years		Repeat Every 10 years
Hepatitis A	Speak to your physician if you need	Speak to your physician if you need	Speak to your physician if you need		
Hepatitis B	Recommended 0,1,6 months	Recommended- High Dose 0,1,2,6 months	Recommended		0,1,2, 6 months
Measles-Mumps-Rubella	1-2 doses (4 weeks apart) *	1-2 doses (4 weeks apart)*	Do not use-Avoid		*Give if Born after 1957, no vaccine, non immune
Chickenpox (Varicella)	2 doses – 4-8 weeks apart	2 doses – 4-8 weeks apart	Do not use-Avoid		
Meningococcal	Speak to your physician if you need	<i>Speak to your physician if you need</i>	Speak to your physician if you need		
Shingles (Shingrix-RZV) For adults over 50 without vaccination	2 doses – 2-6 mos apart	2 doses – 2-6 mos apart	Inactivated virus only		2 doses (2 months apart) Shingrix
RSV	Speak to your physician if you need	Speak to your physician if you need	Speak to your physician if you need		
Travel Vaccines	Speak to your physician if you need	Speak to your physician if you need	Avoid live vaccination		
HPV-Human Papillomavirus	2-3 doses up to age 26	2-3 doses up to age 26	3 doses up to age 26		